

# The Medical Journal of Australia

15 February 1999  
Volume 170 number 4



Established 1914

Journal of the Australian Medical Association

---

## Hospital in the home: a randomised controlled trial

Gideon A Caplan, John A Ward, Nicholas J Brennan, Janis Coconis,  
Neville Board, Ann Brown

**AMPCo**

Australasian Medical Publishing Company Limited, ACN 000 005 854, Level 1, 76 Berry Street, North Sydney, New South Wales 2060, Australia.  
Telephone: (02) 9954 8666 • International +612 9954 8666 • Facsimile (02) 9956 7644 • Email [editorial@ampco.com.au](mailto:editorial@ampco.com.au)

---

*Reprinted from the Medical Journal of Australia • Copyright: Australasian Medical Publishing Company Limited*

---

# Hospital in the home: a randomised controlled trial

Gideon A Caplan, John A Ward, Nicholas J Brennan, Janis Coconis, Neville Board and Ann Brown

Acute care of patients at home is one of the fastest-growing healthcare sectors in the United States<sup>1-3</sup> and is gaining acceptance in many countries.<sup>4</sup> Although there have been randomised controlled trials of patients receiving home versus hospital treatment for deep venous thrombosis,<sup>5</sup> there are few data from trials involving other conditions, or assessing the safety of acute care at home, especially for older patients. It is these patients who occupy an increasing proportion of hospital beds and may derive most benefit from home treatment.<sup>6,7</sup>

Recent studies characterising hospital-associated adverse events<sup>8,9</sup> make home treatment more attractive,<sup>10</sup> especially for elderly patients who are known to suffer a higher incidence of complications in hospital.<sup>11</sup> When treated at home, patients do not have to change their environment or routine, they are not exposed to nosocomial infection, and they do not need to adapt to the sociological culture of the hospital. On the other hand, the incidence and severity of intrinsic complications in older patients can be minimised by round-the-clock supervision by expert staff in hospital. However, there is inadequate evidence to support the popular belief that programs which transfer care to the home result in a worse outcome for the patients.<sup>12,13</sup>

We have conducted a randomised controlled trial of a hospital-in-the-home (HIH) program targeting older patients. We selected conditions which were

## Abstract

**Objectives:** To compare treatment of acute illness at home and in hospital, assessing safety, effect on geriatric complications, and patient/carer satisfaction.

**Design:** Randomised controlled trial.

**Setting:** A tertiary referral hospital affiliated with the University of New South Wales.

**Participants:** 100 patients (69% older than 65 years) with a variety of acute conditions, who were assessed in the emergency department as requiring admission to hospital.

**Interventions:** Patients were allocated at random to be treated by a hospital-in-the-home (HIH) service in their usual residence or to be admitted to hospital.

**Main outcome measures:** Geriatric complications (confusion, falls, urinary incontinence or retention, faecal incontinence or constipation, phlebitis and pressure areas), patient/carer satisfaction, adverse events, and death.

**Results:** There was a lower incidence of confusion (0 v. 20.4% [95% CI, 9.1%–31.7%];  $P=0.0005$ ), urinary complications (incontinence or retention) (2.0% [95% CI, –1.8%, 5.8%] v. 16.3% [95% CI, 6.0%, 26.6%];  $P=0.01$ ), and bowel complications (incontinence or constipation) (0 v. 22.5% [95% CI, 10.7%, 34.1%];  $P=0.0003$ ) among HIH-treated patients. No significant difference in number of adverse events and deaths (to 28 days after discharge) in the two groups was found (although numbers were small). Patient and carer satisfaction was significantly higher in the HIH group.

**Conclusions:** Home treatment appears to provide a safe alternative to hospitalisation for selected patients, and may be preferable for some older patients. We found high levels of both patient and carer satisfaction with home treatment.

MJA 1999; 170: 156-160

amenable to home treatment, but tried to include a variety of diagnostic groups. To accentuate the difference between the two groups, the HIH patients were taken home on the day of presentation to the emergency department, or on the following morning if they presented at night. We studied the safety, efficacy and patient/carer satisfaction of home treatment compared with hospital treatment.

## Methods

### Patients and randomisation

One hundred patients requiring admission to hospital and meeting criteria for study inclusion were randomised to HIH treatment (intervention group) or treatment in hospital (control group). We targeted patients older than 65 years, including those living in nursing homes, but also accepted younger patients. Patients were accepted only if they had been assessed as requiring admission by the relevant medical or surgical team. The medical, surgical and emergency department staff were encouraged to refer patients with acute (pneumonia, urinary tract infections and cellulitis) and subacute (endocarditis and osteomyelitis) infections requiring treatment with intravenous antibiotics, deep venous thrombosis (DVT), minor cerebrovascular accidents (not affecting mobility or swallowing), and cardiac failure.

## For editorial comment, see page 148

### Post Acute Care Services, Prince of Wales Hospital, Sydney, NSW.

Gideon A Caplan, MB BS, FRACP, Director; and Lecturer, University of New South Wales,

Sydney, NSW; Nicholas J Brennan, MB BS, FRACP, Senior Registrar;

Janis Coconis, RN, Clinical Nurse Specialist; Neville Board, RN, BA, Research Assistant;

Ann Brown, RN, Nurse Manager.

### Community Health Services and Programs, South Eastern Sydney Area Health Service, Sydney, NSW.

John A Ward, MB BS, MSc, FRACP, Community Geriatrician; and Lecturer, University of New South Wales, Sydney, NSW.

Reprints: Dr G A Caplan, Post Acute Care Services, Prince of Wales Hospital, Randwick, NSW 2031.

Email: g.caplan@unsw.edu.au

Patients were excluded from the study

- if they had evidence of shock (systolic blood pressure <100 mmHg);
- if they required oxygen (PaO<sub>2</sub> <60 mmHg);
- if they were judged too unwell by the study team;
- if they had no available carer;
- if they lived outside the local area; or
- if their home was unsuitable for home treatment (lack of running water, electricity or an inside toilet, concerns about safety, or dangerous pets).

If any doubts about home suitability arose during the emergency department assessment, a study nurse visited the home before randomisation.

After informed consent was obtained from the patient (written) and the patient's carer (verbal), randomisation (stratified according to whether the patient lived at home or in a nursing home, or had DVT) was achieved by computer-generated random numbers coded into sealed envelopes.

The study protocol required that patients in the HIH group be discharged from hospital within 24 hours of diagnosis. HIH patients receiving intravenous antibiotics were given the first dose in hospital. The study protocol was approved by the hospital's Research Ethics Committee.

**Data collection**

For all study patients, study nurses completed a Barthel Index of Activities of Daily Living,<sup>14</sup> a modified Instrumental Activities of Daily Living Index<sup>15</sup> and a Mental Status Questionnaire<sup>16</sup> on admission and discharge. Baseline demographic data on current medical diagnoses, a detailed social history, physical function, medications and allergies were also recorded.

**Hospital treatment**

Hospital patients (control group) were admitted under the appropriate physician or surgeon of the day and treated in accordance with standard regimens without the intervention of the study team. The hospital team treating the patient was notified that the patient had been included in the trial.

**HIH treatment**

HIH patients (intervention group) were treated according to the presenting diagnosis by the hospital community outreach team. The range of treatments

in the study protocol included administration of parenteral antibiotics and other medications, and blood transfusions. Infections were generally treated with once-daily intravenous antibiotics such as ceftriaxone, gentamicin or vancomycin, as appropriate, according to the result of bacteriological tests (if positive) or by diagnosis. Intravenous access was usually via cannulas inserted by study nurses. Owing to safety concerns, some patients with dementia did not have cannulas left *in situ*, and were cannulated daily with butterfly cannulas. Patients requiring long term treatment or with difficult venous access were treated via a peripherally inserted central cannula. Patients with DVT were treated with daily subcutaneous enoxaparin injections (1.5 mg/kg) and oral warfarin until their international normalised ratio was 2.0 or above.

**GP involvement**

Before the study, we conducted an educational program consisting of an evening lecture and a question-and-answer session for the local division of

general practice. Study nurses also visited general practitioners (GPs) and nursing homes to explain the program. Our preference was for the patient's own GP to be the primary medical manager, and we therefore sought the agreement of the patient's GP before entering the patient in the trial. If GPs declined to participate, the hospital provided medical support.

**Follow-up**

After discharge, an unmarked satisfaction survey based on the principles outlined by Draper and Hill,<sup>17</sup> colour coded to differentiate between responses from HIH and hospital treatment, was sent to patients, carers and GPs. This was followed by a phone call to encourage response. Patients were asked "How would you rate your treatment overall?" and offered a four-point scale of answers: excellent (1), good (2), fair (3), and poor (4). A similar question about satisfaction with the patient's care was asked of the carer and GP. Patients were recontacted by telephone at one and six

**1: Characteristics of the 100 patients entered into the trial. Values are means with 95% confidence intervals unless stated otherwise**

	Hospital in the home (n=51)	Hospital (n=49)
Age in years (median, range)	73, 17-111	79, 22-97
Number (%) of patients ≥65 years	36 (70.6%)	33 (67.3%)
Male:female ratio	21:30	24:25
Time in emergency department (h)	7.8 (5.9, 9.7)	12.2 (10.1, 14.3)
Length of stay (days)	10.1 (7.7, 12.5)	7.4 (6.0, 8.8)
Function measures		
Barthel Index (maximum score = 20)	15.2 (13.3, 17.1)	14.8 (12.8, 16.8)
Instrumental Activities of Daily Living (maximum score = 12)	6.8 (5.5, 8.1)	6.2 (4.7, 7.7)
Mental Status Questionnaire (maximum score = 10)	7.1 (6.1, 8.1)	6.9 (5.8, 8.0)
Creatinine level (mmol/L)	0.10 (0.08, 0.12)	0.11 (0.10, 0.12)
Type of residence (number [%] patients)		
Home	37 (72.5%)	33 (67.3%)
Nursing home	12 (23.5%)	13 (26.5%)
Hostel	2 (3.9%)	3 (6.1%)
Intravenous antibiotics group	n=37	n=34
White cell count (x10 <sup>9</sup> /L)	10.2 (8.6, 11.8)	11.8 (10.0, 13.6)
Temperature (°C)	37.1 (36.9, 37.3)	37.5 (37.3, 37.7)
Number (%) patients with positive cultures	13 (35.1%)	12 (35.3%)

\* Differences between patients treated at home and those treated in hospital are not significant, except for time spent in the emergency department (P < 0.003).

pital-treated group, respectively, were recorded as having an unplanned readmission within 28 days of discharge.

There was one death in each of the HIH- and hospital-treated groups during the admission. Total deaths up to 28 days after discharge were three in the HIH group and four in the hospital-treated group, and at six months after discharge six and seven deaths, respectively, had occurred. There were no significant differences between the two groups in these outcomes.

#### Patient, carer and GP satisfaction

The response rates for the satisfaction surveys were (HIH v. control): patients 78% v. 40%, carers 55% v. 27%, and GPs 63% v. 37%. There was significantly greater satisfaction with overall HIH treatment among both patients ( $P < 0.0001$ ) and their carers ( $P = 0.0001$ ). Despite the increased workload and despite having to deal with more severely ill patients than usual, GPs were equally satisfied by home and hospital treatment (Table 4).

#### Discussion

In our study, treatment of a range of acute infections and other diseases at home was as safe as treatment in hospital. Previous studies which claimed to have demonstrated safety documented so few complications that risk appeared to have been eliminated by patient selection.<sup>1-3</sup> Our study group's 7% mortality up to 28 days after discharge (compared with zero mortality in previous studies<sup>1-3</sup>) establishes our patients' illness severity.

The most significant finding resulting from our inclusion of geriatric complications was the higher incidence of confusion in hospital-treated patients. The role of the hospital environment in the aetiology of delirium has long been recognised,<sup>21</sup> although this is not universally accepted.<sup>22</sup> For some of the patients in our study recorded as having confusion, all of the features of delirium according to the *Diagnostic and statistical manual of mental disorders* (DSM-IV)<sup>23</sup> were not present. Our aim was to be as

4: Satisfaction survey — mean responses and 95% confidence intervals

	Hospital in the home (n=51)	Hospital (n=49)	P
Patient			
Mean score‡	1.1	2.0	<0.0001
95% CI	1.0, 1.2	1.7, 2.3	
Carer			
Mean score‡	1.1	1.9	0.0001
95% CI	1.0, 1.2	1.4, 2.4	
General practitioner			
Mean score‡	1.7	1.8	ns
95% CI	1.4, 2.0	1.4, 2.2	

\*Satisfaction was rated on a 4-point scale: 1 = Excellent; 2 = Good; 3 = Fair; 4 = Poor. na = not applicable.

inclusive as possible of geriatric complications, so that if a problem was noted by ward staff or carers it was included even if study personnel had not observed it. These complications were not always serious clinically and often did not have a great impact, but we valued the perspective of patients, families and hospital staff. Similarly, the occurrence of urinary and bowel complications reflects the difficulty of caring for frail elderly patients<sup>24</sup> (especially those from nursing homes) in an acute hospital ward. Patients with mobility or communication problems require carers attuned to their routines and signals to maintain continence.

Because of small numbers our study was limited in its power to draw conclusions about the difference between HIH and hospital treatment as far as adverse events and deaths were concerned. As there have been no previous studies addressing this question, it was not possible to estimate the size of the expected difference. Another limitation of our study was that, although we did not directly inform the adverse event reviewers about which patients were in each group, we were not able to completely blind them to the status of each group because of stylistic differences between the patient records of the hospital- and the HIH-treated patients.

Few similar comparative studies have been published. As reports and discussions entitled "home hospitalisation" or "hospital at home", for example, may include few or no acute patients, the home treatment of acute illness is obscured in the medical literature.<sup>25,26</sup> Our study included only patients who met the criterion of requiring admission

to hospital because of illness. By bringing the hospital care to the patient's home, we substituted completely for hospital facilities. There is a need for more studies of patients with illness of this severity treated at home.

A number of factors may explain the greater satisfaction with treatment at home. In their own homes, patients are in their familiar environment, their privacy is protected, their sleep less interrupted, and they can eat their usual food.<sup>27,28</sup> As an example, for

an elderly patient with limited cognitive function and/or decreased visual acuity, finding the bathroom at night in hospital may be impossible. Moreover, elderly patients may be unable to get out of bed because of rails on the sides of the bed. These difficulties may have contributed to the higher incidence of incontinence in our hospital-treated group. At home, patients are more active participants, often partners, in their care. All these factors invariably translate into greater satisfaction with home treatment.<sup>29</sup>

Frail, older patients — because of their atypical disease presentations, more rapid deterioration when ill and greater prevalence of disease — need hospital-level services more frequently than younger patients. However, admission to hospital, like any medical intervention, has side effects as well as benefits.<sup>8</sup> Elderly patients have more gain from many treatments (eg, thrombolytic therapy) because of their higher underlying mortality rates,<sup>30</sup> but they also suffer from a much greater occurrence of iatrogenic complications.<sup>10</sup> Although many of these complications may be explained by an interaction between their underlying frailty and their disease process,<sup>8</sup> we were able to detect a measurable effect attributable to a change of environment.

The results of this study do not imply that older patients should never be admitted to hospital. On the contrary, the elderly as a group have more need for the technical expertise of the acute hospital than younger people. However, if such technical expertise can be delivered at home, this alternative should at least be evaluated. On the other hand, our study does not imply that treatment

which can be delivered at home should always be delivered at home. Successful home treatment depends on careful patient and home selection, as the necessary minimum level of sanitation, facilities and support from carers is not universally available.<sup>4</sup> However, where it is feasible, we have demonstrated that home treatment of acute illness, even for frail elderly patients, can be effective, safe and more satisfying for the patient and their families.

### Acknowledgements

We would like to acknowledge the invaluable assistance of the staff of Post Acute Care Services and the Emergency Department at Prince of Wales Hospital and Ms Bernie Harrison and Dr Ross Wilson from QARNS at Royal North Shore Hospital.

This study was supported by a grant from the Commonwealth Department of Health and Family Services - Hospital Access (National Projects) Program.

### References

- Antoniskis A, Anderson BC, Von Volkinburg EJ, et al. Feasibility of outpatient self-administration of parenteral antibiotics. *West J Med* 1978; 128: 203-206.
- Kind AC, Williams DN, Persons G, et al. Intravenous antibiotic therapy at home. *Arch Intern Med* 1979; 139: 413-415.
- Stiver HG, Telford GO, Mossey JM, et al. Intravenous antibiotic therapy at home. *Ann Intern Med* 1978; 89: 690-693.
- Tice AD. Outpatient parenteral antibiotic therapy in different countries. *Int J Infect Dis* 1996; 1: 102-106.
- Levine M, Gent M, Hirsh J, et al. A comparison of low-molecular-weight heparin administered primarily at home with unfractionated heparin administered in the hospital for proximal deep-vein thrombosis. *N Engl J Med* 1996; 334: 677-681.
- Balinsky W, Nesbitt S. Cost-effectiveness of outpatient parenteral antibiotics: a review of the literature. *Am J Med* 1989; 87: 301-305.
- Poretz DM, Eron LJ, Goldenberg RL, et al. Intravenous antibiotic therapy in an outpatient setting. *JAMA* 1982; 248: 336-339.
- Leape LL, Brennan TA, Laird N, et al. The nature of adverse events in hospitalised patients: results of the Harvard Medical Practice Study II. *N Engl J Med* 1991; 324: 377-384.
- Wilson RM, Runciman WB, Gibberd RW, et al. The quality in Australian health care study. *Med J Aust* 1995; 163: 458-471.
- Caplan GA, Brown A. Post acute care: can hospitals do better with less? *Aust Health Rev* 1997; 20: 43-54.
- Fretwell MD. Acute hospital care for frail older patients. In: Hazzard WR, Andres R, Bierman EL, Blass JP editors. *Principles of geriatric medicine and gerontology*, 2nd ed. New York: McGraw-Hill, 1990: 247-253.
- Maxwell RJ. Why rationing is on the agenda. *Br Med Bull* 1995; 51: 761-768.
- Larkins RG, Martin TJ, Johnston CI. The boundaryless hospital — a commentary. *Aust N Z J Med* 1995; 25: 169-170.
- Mahoney FI, Barthel DW. Functional evaluation: the Barthel index. *Md Med J* 1965; 14: 61-65.
- Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969; 9: 179-186.
- Pfeiffer E. A short portable mental status questionnaire for the assessment of organic brain deficit in elderly patients. *J Am Geriatr Soc* 1975; 23: 433-441.
- Draper M, Hill S. The role of patient satisfaction surveys in a national approach to hospital quality management. Canberra: AGPS, 1996.
- Hazzard WR, Andres R, Bierman EL, Blass JP editors. *Principles of geriatric medicine and gerontology*. 2nd ed. Section 4 — geriatric syndromes and special problems. New York: McGraw-Hill, 1990: 1055-1222.
- SPSS for Windows [computer program], version 6.0. Chicago: SPSS Inc, 1993.
- Dean AG, Dean JA, Burton AH, Dicker RC. Epi-Info [computer program], version 5. Stone Mountain, Ga: USD Inc, 1990.
- Wahl CW, Golden JS, Liston EH, et al. Toxic and functional psychoses: diagnosis and treatment in a medical setting. *Ann Intern Med* 1967; 66: 989-1007.
- Lipowski ZJ. Delirium and impaired consciousness. In: Evan JG, Williams TF, editors. *Oxford textbook of geriatric medicine*. Oxford: Oxford University Press, 1992: 490-495.
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (DSM-IV)*. Washington, DC: APA, 1994.
- Caplan GA, Brown A, Croker WD, Doolan J. Risk of admission within four weeks of discharge of elderly from the emergency department — the DEED study. *Age Ageing*. In press.
- Stressman J, Ginsberg G, Hammerman-Rozenberg R, et al. Decreased hospital utilization by older adults attributable to a home hospitalization program. *J Am Geriatr Soc* 1996; 44: 591-598.
- Donald IP, Baldwin RN, Bannerjee M. Gloucester hospital-at-home: a randomized controlled trial. *Age Ageing* 1995; 24: 434-439.
- Consumer reports of hospital experiences. In: Draper M, Hill S, editors. *The role of patient satisfaction surveys in a national approach to hospital quality management*. Canberra: AGPS, 1996: 12-20.
- Cruse PJE, Foord R. The epidemiology of wound infection: a 10-year prospective study of 62,939 wounds. *Surg Clin North Am* 1980; 60: 27-40.
- Brown RB. Selecting the patient. *Hosp Pract* 1993; 28(Suppl 1): 11-15.
- Fibrinolytic Therapy Trialists' (FTT) Collaborative Group. Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet* 1994; 343: 311-322.

(Received 17 Jun, accepted 22 Oct, 1998)

□